

AD-A043 490

FLORIDA STATE UNIV TALLAHASSEE DEPT OF STATISTICS
AN INEQUALITY FOR CONVEX FUNCTIONS INVOLVING G-MAJORIZATION.(U)

F/6 12/1

JUL 77 R V LEON, F PROSCHAN

AF-AFOSR-2581-74

UNCLASSIFIED

FSU-STATISTICS-M412

AFOSR-TR-77-1148

NL

| OF |
AD
A043490



END

DATE

FILMED

9-77

DDC

ADA043490

An Inequality for Convex Functions Involving
G-Majorization

by

Ramón V. León¹ and Frank Proschan²
The Florida State University

FSU Statistics Report M412
AFOSR Technical Report No. 73

July, 1977
The Florida State University
Department of Statistics
Tallahassee, Florida 32306

DDC
RECEIVED
AUG 29 1977
B

¹Research supported by National Institute of Environmental Health Sciences under Grant 1 T32 ES07011 at Florida State University.

²Research supported by the Air Force Office of Scientific Research, AFSC, USAF, under Grant AFOSR-74-2581.

The United States Government is authorized to reproduce and distribute reprints for government purposes.

AMS Numbers: 26A86
26A48

DISTRIBUTION STATEMENT A
Approved for public release
Distribution Unlimited

AD NO. —
DDC FILE COPY

3

R

ALOSR-7-1148

064840VADA

An Inquiry to Formosa Province Involving
G-400000000

by

Research, Design and Production
Department State Division

Administrative Report
Technical Report

AIR FORCE OFFICE OF SCIENTIFIC RESEARCH (AFSC)
NOTICE OF TRANSMITTAL TO DDC
This technical report has been reviewed and is
approved for public release IAW AFR 190-12 (7b).
Distribution is unlimited.
A. D. BLOSE
Technical Information Officer

000 LKE COPY

An Inequality for Convex Functions Involving
G-Majorization

by

by Ramón V. León and Frank Proschan
The Florida State University

ABSTRACT

In this paper we derive a simple inequality involving expectations of convex functions and the notion of G-majorization. The result extends a similar inequality of Marshall and Proschan (1965), J. Math. Anal. Applic. Useful applications of the more general inequality are presented.

ACCESSION for	
NTIS	White Section <input checked="" type="checkbox"/>
DDC	Buff Section <input type="checkbox"/>
UNANNOUNCED	<input type="checkbox"/>
JUSTIFICATION	
BY	
DISTRIBUTION/AVAILABILITY CODES	
Dist. AVAIL. and/or SPECIAL	
A	

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 18 AFOSR-TR-77-1148	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER 9 Technical rep.
4. TITLE (and Subtitle) 6 AN INEQUALITY FOR CONVEX FUNCTIONS INVOLVING G-MAJORIZATION.	5. TYPE OF REPORT & PERIOD COVERED	
7. AUTHOR(s) 10 Ramon V. Leon Frank/Proschan	6. PERFORMING ORG. REPORT NUMBER FSU Stat. Report No M412 L	
9. PERFORMING ORGANIZATION NAME AND ADDRESS The Florida State University Department of Statistics Tallahassee, FL 32306	8. CONTRACT OR GRANT NUMBER(s) 15 AF-AFOSR 24-2581-74	
11. CONTROLLING OFFICE NAME AND ADDRESS Air Force Office of Scientific Research/NM Bolling AFB, Washington, DC 20332	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 61102F 16 2304 AS 17 A5	
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) 14 FSU-Statistics-M412	12. REPORT DATE 11 Jul 1977	
	13. NUMBER OF PAGES 5 129 p.	
	15. SECURITY CLASS. (for this report) UNCLASSIFIED	
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) majorization, G-majorization, convex functions, inequalities, permutations, sign changes, groups, partial ordering		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) In this paper we derive a simple inequality involving expectations of convex function and the notion of G-majorization. The result extends a similar inequality of Marshall and Proschan (1965), J. Math. Anal. <u>Applic.</u> Useful applications of the more general inequality are presented. 400277 <i>Ine</i>		

An Inequality for Convex Functions Involving
G-Majorization

Ramón V. León and Frank Proschan
The Florida State University

In this note we derive a simple inequality involving expectations of convex functions and the notion of G-majorization. The result extends a similar inequality of Marshall and Proschan (1965) involving majorization. A number of useful applications of the more general inequality are then presented.

Let G be a group of matrices (linear transformations) acting on R^n . A vector $a = (a_1, \dots, a_n)$ is said to G-majorize a vector $b = (b_1, \dots, b_n)$, written $a \overset{G}{\succeq} b$, if b is in the convex hull of the G -orbit of a . If $G = P_n$, the group of permutation matrices, G -majorization coincides with majorization (see Eaton and Perlman, 1976). A random vector $X = (X_1, \dots, X_n)$ is said to be G-invariant if X is stochastically equal to gX for all $g \in G$. When $G = P_n$, we say that X_1, \dots, X_n are exchangeable random variables. For vectors a and b , let $a \cdot b \stackrel{\text{def}}{=} (a_1 b_1, \dots, a_n b_n)$.

Theorem 1. Let G be a finite group such that for all $g \in G$ there exist h and $k \in G$ for which $h(ga \cdot b) = a \cdot kb$ for all vectors a and b . Let X be a G -invariant random vector, ϕ a continuous, convex, G -invariant function and $a \overset{G}{\succeq} b$. Then

$$(1) \quad E\phi(a \cdot X) \geq E\phi(b \cdot X).$$

Moreover, if ϕ is strictly convex, equality holds only when $a = gb$ for some $g \in G$, or when the X_i are all zero with probability one.

Proof. Let $G = \{g_i\}_{i=1}^m$. Then we may write $b = \sum_{j=1}^m \alpha_j g_j a$, where each $\alpha_j \geq 0$ and $\sum_{j=1}^m \alpha_j = 1$. It follows that $E\phi(b \cdot X) = E\phi([\sum_{j=1}^m \alpha_j g_j a] \cdot X) =$

$E\phi(\sum_{j=1}^m \alpha_j [g_j a \cdot X]) \leq \sum_{j=1}^m \alpha_j E\phi(g_j a \cdot X)$. For each j , let h_j and k_j be the elements of G for which $h_j(g_j a \cdot X) = a \cdot k_j X$. Then $E\phi(g_j a \cdot X) = E\phi(h_j(g_j a \cdot X))$ [by the G -invariance of ϕ] $= E\phi(a \cdot k_j X) = E\phi(a \cdot X)$ [by the G -invariance of X]. Thus $E\phi(b \cdot X) \leq \sum_{j=1}^m \alpha_j E\phi(a \cdot X) = E\phi(a \cdot X)$.

In case ϕ is strictly convex, it is clear from the above proof that equality holds only if for some $g \in G$, $b \cdot X = ga \cdot X$ with probability one. \square

Remark 1. If $G = P_n$, then for all $g \in G$ and vectors a and b , $g^{-1}(ga \cdot b) = a \cdot g^{-1}b$. Therefore in this special case the hypothesis of Theorem 1 is satisfied. It follows that the main result of Marshall and Proschan (1965) involving majorization is a special case of Theorem 1.

Remark 2. Other groups of interest for which the hypothesis of the theorem is satisfied are: (a) The group G_1 of sign changes and (b) the group G_2 of permutations and sign changes, as is readily verified.

Remark 3. Note that in G_2 , $g^{-1}(ga \cdot b) = a \cdot g^{-1}b$ for all $g \in G_2$. For example if $g = \begin{bmatrix} 0 & -1 \\ -1 & 0 \end{bmatrix}$ then $\begin{bmatrix} 0 & -1 \\ -1 & 0 \end{bmatrix} \left(\begin{bmatrix} 0 & -1 \\ -1 & 0 \end{bmatrix} a \cdot b \right) = a \cdot \left(\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} b \right)$. So in G_2 the milder requirement is needed that for all $g \in G$ there exist h and $k \in G$ for which $h(ga \cdot b) = a \cdot kb$. Also note that this condition is not satisfied for some groups. For example, if $G = \left\{ \begin{bmatrix} 0 & -1 \\ -1 & 0 \end{bmatrix}, I \right\}$ and $g = \begin{bmatrix} 0 & -1 \\ -1 & 0 \end{bmatrix}$, then clearly there do not exist h and k for which $h(ga \cdot b) = a \cdot kb$.

Remark 4. Let $G = G_2$ and let vectors a and b have all components non-negative. Then $a \stackrel{G}{\leq} b$ if and only if a is weakly majorized by b . (See Marshall, Walkup and Wets (1967) for the definition of weak majorization.)

Similarly, let $G = G_1$, and let vectors a and b have all components

nonnegative. Then $\underline{a} \stackrel{G}{\leq} \underline{b}$ if and only if $a_i \leq b_i$ for $i = 1, 2, \dots, n$. It follows that Theorem 1 yields results concerning weak majorization and the usual partial ordering of the plane. (See also Remark 8.)

Remark 5. For comments on a converse to Theorem 1, see Remark 3 of Marshall and Proschan (1965). Also see Remark 4 of that paper for a counterexample showing that the conclusion of Theorem 1 need not necessarily hold when we weaken the hypothesis to require ϕ to be only continuous and isotone with respect to the G-majorization ordering, i.e., G-monotone. (A G-invariant convex function is necessarily G-monotone). However, by using a path lemma of Eaton and Perlman (1976), it is possible to show that if G is a reflection group, then Theorem 1 holds when ϕ is merely continuous and convex along all the line segments joining \underline{a} with $g\underline{a}$ for all $g \in G$. (See Eaton and Perlman (1976) for the definition of a reflection group.) This is consistent with Remark 4 of Marshall and Proschan (1965). Thus if $G = G_2$ we need only require that ϕ , considered as a function of a specified pair of coordinates with all other coordinates held fixed, be convex. Note that this condition on ϕ is the same as that in Remark 4 of Marshall and Proschan (1965). Similarly, if $G = G_1$, we need only require that ϕ , considered as a function of a specified coordinate with all other coordinates held fixed, be convex.

Corollary 1. Let $G = G_2$. Let $X(\sigma_1), \dots, X(\sigma_n)$ be independent random variables, where $X(\sigma_i)$ is normally distributed with mean zero and standard deviation σ_i . Let ϕ be continuous, convex, and G-invariant, and $(\sigma_1, \dots, \sigma_n) \stackrel{G}{\geq} (\sigma_1', \dots, \sigma_n')$. Then

$$(2) \quad E\phi(X(\sigma_1), \dots, X(\sigma_n)) \geq E\phi(X(\sigma_1'), \dots, X(\sigma_n')).$$

Proof. Let Y_1, \dots, Y_n be independently distributed standard normal random

variables. Then $E\phi(\sigma_1 Y_1, \dots, \sigma_n Y_n) \geq E\phi(\sigma_1 Y_1, \dots, \sigma_n Y_n)$ by Theorem 1.

Since $\sigma_i Y_i$ and $X(\sigma_i)$ have the same distribution, the result follows. \square

Remark 6. Similar results are true when G is P_n or G_1 .

Remark 7. Note that the only property of $X(\sigma_i)$ used in the proof of Corollary 1 is that $X(\sigma_i)$ and $\sigma_i Y$ have the same distribution where Y is a random variable distributed symmetrically about zero. Thus, for example, Corollary 1 is still true when $X(\sigma_i)$ is uniformly distributed on the interval $(-\sigma_i, \sigma_i)$. For simplicity, Corollary 1 is stated for the special case $X(\sigma_i)$ is normal.

Remark 8. Note that since $\overset{P_n}{a} \leq \overset{G_1}{b}$ or $\overset{G_1}{a} \leq \overset{G_2}{b}$ implies $\overset{G_2}{a} \leq \overset{G_2}{b}$, (2) holds when $\overset{P_n}{a} \leq \overset{G_1}{b}$ or $\overset{G_1}{a} \leq \overset{G_2}{b}$. (A similar remark applies whenever the G_2 ordering holds.)

Corollary 2. Let ϕ be continuous, convex, and invariant under permutations and sign changes, and let $\overset{G_2}{a} \geq \overset{G_2}{b}$. Then

(3) $\sum^* \phi(a_1 x_1, \dots, a_n x_n) \geq \sum^* \phi(b_1 x_1, \dots, b_n x_n)$ where \sum^* denotes summation over all sign changes and permutations of the x_i .

Proof. (3) is an immediate consequence of (1), where $P\{(x_1, \dots, x_n) = ((-1)^{\sigma_1} x_{i_1}, \dots, (-1)^{\sigma_n} x_{i_n})\} = \frac{1}{2^n n!}$, where (i_1, \dots, i_n) is a permutation of $(1, 2, \dots, n)$ and $\sigma_i = 0$ or 1 for $i=1, \dots, n$.

Remark 9. Corollary 2 is a variation of Muirhead's Theorem. (See Hardy, Littlewood, and Pólya, 1952, pp. 44-48.)

Remark 10. For other possible applications yielding inequalities, see Marshall and Proschan (1965).

REFERENCES

- Conlon, J. C., León, R., Proschan, F., and Sethuraman, J. (1977). G-Ordered Functions in Statistics. AFOSR Report 71, Dept. of Statistics, Florida State University.
- Eaton, M. L. and Perlman, M. D. (1976). Reflection Groups, Generalized Schur Functions, and the Geometry of Majorization. University of Chicago Statistics Report No. 28.
- Hardy, G. H., Littlewood, J. E., Pólya, G. (1952). Inequalities. Cambridge University Press.
- Jogdeo, K. (1977). Association and Probability Inequalities. Ann. Statist. 5 495-504.
- Marshall, A. W. and Proschan, F. (1965). An Inequality for Convex Functions Involving Majorization. J. Math. Anal. Applic. 12 87-90.
- Marshall, A. W., Olkin, D. W. and Wets, R. (1967). Order-preserving Functions; Applications to Majorization and Order Statistics. Pacific J. Math. 23 584.











